



Ebola Virus Disease (EVD) and COVID-19 Screening Form for Health Care Facilities

SECONDARY SCREENING to risk stratify for EVD

ASK EACH PATIENT THAT HAS SCREENED "YES" TO PRIMARY SCREENING QUESTIONS, ASK THE FOLLOWING QUESTIONS:

Have you experienced **ANY** of the following recently?

Exposure

- Attended funeral of a family member or friend
- Contact with sick or dead animal
- Attended a party/social gathering/religious events during EVD epidemic
- Sexual contact with male EVD survivors

Symptoms

- Difficulty swallowing
- Skin rash

Patient responds **NO** to **ALL** criteria

↓ NO ↓

✓ Go to question 3: Assess for COVID-19

Patient responds **YES** to **ANY** of these criteria

↓ YES ↓

- ! Place patient in isolation area. Ideally, EOC authorities/rapid response team will be notified immediately and arrive quickly to either assess and/or take suspected EVD case to further management.
- If patient must wait at PHC for some time, trained clinicians should don PPE for suspected EVD case using "buddy" system, conduct exam/intake after notifying EOC authorities.
- If possible, trained clinical should test patient for EVD using rapid test if available.
- If test is positive for EVD, patient should be kept in isolation and provided supportive care, and the local or EOC authorities should be notified to arrange transport to an Ebola Treatment Center (ETU).
- If possible, patients with suspected EVD should be kept in a separate space from those with suspected COVID-19. If separation is not possible, contact your local authorities for guidance.
- Consider testing for malaria as well if resources available.
- If test is negative for EVD, go to tertiary screening.

TERTIARY SCREENING to assess for COVID-19

ASK EACH PATIENT THAT SCREENED "YES" TO SECONDARY SCREENING QUESTIONS, ASK THE FOLLOWING QUESTIONS:

Have you experienced **ANY** of the following recently?

Symptoms

- Cough
- Shortness of breath
- Sore throat
- Loss of smell/taste
- Shortness of breath
- Congestion or running nose

Patient responds **NO** to **ALL** criteria

↓ NO ↓

✓ Assess for other conditions /permit entry into HCF

Patient responds **YES** to **ANY** of these criteria

↓ YES ↓

- ! Place patient in isolation space for potential COVID-19 suspects or keep patient outdoors, separated from others by at least 1-meter if possible.
- Trained clinician should maintain at least 1-meter distance from patient while performing screening/exam. If not possible to maintain physical distance from patient, the clinician should don PPE for COVID-19 (mask, gloves, goggles/face shield, gown (if available)).
- If possible, test for COVID-19 with rapid test.
- If test is positive, maintain patient separated/isolated; COVID-19 patient pathway/clinical management is active.
- Notify local authorities/surveillance teams of potential COVID-19 case.